arXiv:1211.1947v1 [hep-ph] 8 Nov 2012

# Soft and Collinear Factorization and Transverse Momentum Dependent Parton Distribution Functions

Miguel G. Echevarría,[1,*] Ahmad Idilbi,[2,†] and Ignazio Scimemi[1,‡]
[1]*Departamento de Física Teórica II, Universidad Complutense de Madrid (UCM), 28040 Madrid, Spain*
[2]*European Centre for Theoretical Studies in Nuclear Physics and Related Areas (ECT*),*
*Villa Tambosi, Strada delle Tabarelle 286, I-38123, Villazzano, Trento, Italy*

We consider the factorization of hadronic physical observables with transverse momentum dependence when two collinear sectors are involved and which interact through emission of soft gluons. By proving a novel property of the relevant soft function as a geometric mean of two "hemisphere" soft functions, one can combine those functions with the two collinear ones to obtain a modified factorization theorem in which all quantities are well-defined and each "generalized" collinear quantity belongs to distinct region in rapidity space. Those quantities will be identified as the "transverse momentum dependent parton distribution functions". They are free from all rapidity divergences, including the mixed ultraviolet/non-ultraviolet ones, their evolution is governed by a homogeneous differential equation and no auxiliary parameters are needed to resum large logarithms.

**Introduction:** Transverse momentum dependent (TMD) factorization theorems are essential theoretical tools in modern high-energy colliders. High-energy scattering reactions such as semi-inclusive deep-inelastic scattering (SIDIS), the $q_T$-dependent spectrum for Drell-Yan (DY) heavy lepton pair production or the analogous gluon-gluon fusion process through proton-proton collisions at the LHC, all ask for a TMD factorization theorems to be established before carrying out phenomenology. Such theorems have mainly dual purpose. First is to disentangle long and short distance physics while keeping the TMD manifest. This serves to consider $q_T$-dependent spectrums in which the product(s) of high-energy reactions are produced (or tagged) with a non-vanishing transverse momentum. Those events constitute a major bulk of the LHC data. Secondly, when the spin is invoked with polarized targets and/or products then the nucleon's three-dimensional structure as well as its momentum and spin distributions can be unraveled to a maximal extent.

Establishing such TMD factorization theorems is much more complex than the collinear counterparts (where all the TMD is integrated over). The complexity arises since, in certain kinematical regimes, the relevant modes that capture the long-distance hadronic physics, namely soft and collinear [1] (e.g., [3]), have the same invariant mass but differ only in their relative rapidities. In such instances the soft and collinear modes can mix under boosts (see the discussion in Sec. VII of [4] and references therein), thus the scale factorization becomes a subtle issue.

The TMD parton distribution functions (TMDPDFs) where introduced by Collins and Soper [5, 6]. Later on Ji, Ma and Yuan [7] re-defined the TMDPDF by subtracting a complete soft function from the bi-local collinear quark field operator. Since then several works have emerged [3, 8–14] with seemingly contradictory conclusions. In some places it was even concluded that the notion of "TMDPDF" simply cannot be defined at the operator level contrary to the integrated Feynman PDF.

Below we argue that only when the soft function is partitioned –in rapidity space– in a correct way and combined with the collinear matrix elements, one can then separate modes with positive and negative rapidities, thus obtaining a "two-hemisphere"-like picture. As a consequence, we would be able to define TMDPDFs (whether polarized or unpolarized) which do not suffer from any bad features [2]. Mainly they will be free from all rapidity divergences including mixed ultraviolet (UV)/non-ultraviolet (nUV) ones. Those divergences pose a serious problem to the renormalizability of such physical objects. Our approach is that such mixed UV/nUV divergences should not exist, right from the start and before renormalization, in the proper definition of a TMDPDF. Also, and not less importantly, is the ability to recover the integrated PDF from the newly defined TMDPDF without reference to any particular choice of regulators or prescriptions.

---

[1] Depending on the virtuality $p^2$ of the modes they are either described by SCET-II ($p^2 \sim \Lambda^2_{\rm QCD}$) [1] or SCET-$q_T$ ($p^2 \sim q_T^2$) [2].

[2] To best of our knowledge there is not any single treatment so far in which the mixed UV/nUV divergences cancel *regardless* of regulators or prescriptions.

**Outline of the Problem:** Let us consider the DY kinematics in the center-of-mass frame where the momenta of the two incoming partons initiating the hard reaction are $p$ and $\bar{p}$. A general vector $v$ is decomposed as: $v^\mu = \bar{n}\cdot v\frac{n^\mu}{2} + n\cdot v\frac{\bar{n}^\mu}{2} + v_\perp = v^+\frac{n^\mu}{2} + v^-\frac{\bar{n}^\mu}{2} + v_\perp$, where $n = (1,0,0,1)$, $\bar{n} = (1,0,0,-1)$. Then, $p = (Q,0,\vec{0}_\perp)$ and $\bar{p} = (0,Q,\vec{0}_\perp)$ where $Q$ is the virtual photon mass. The relevant modes that contribute to the DY $q_T$-spectrum are collinear ($k \sim Q(1,\lambda^2,\lambda)$), anti-collinear ($k \sim Q(\lambda^2,1,\lambda)$) and soft ($k \sim Q(\lambda,\lambda,\lambda)$), where $\lambda \sim q_T/Q$ is small.

As mentioned earlier the soft modes can become $n,\bar{n}$-collinear under boosts and vice versa. To separate the modes one needs to break the "soft-collinear" boost invariance by introducing rapidity cuts which also serve, in perturbation theory, to regularize rapidity divergences, i.e., when $y \equiv \frac{1}{2}\ln|k^+/k^-| \to \pm\infty$.


$k^+$
$\eta^+$
$\eta^-$
$k^+ = k^-$
$\tilde{J}_n^{(0)}(\eta^-)$
$\sqrt{\tilde{S}(\eta^-,\eta^-)}$
$\eta^-$
$\eta^+$
$\sqrt{\tilde{S}(\eta^+,\eta^+)}$
$B1$
$B2$
$A$
$\tilde{J}_{\bar{n}}^{(0)}(\eta^+)$
$C$
$k^-$

FIG. 1: *Relevant modes for the factorization of DY $q_T$-spectrum. Regions A and C are the pure collinear modes in the $n$ and $\bar{n}$ directions, respectively, and B1+B2 are the soft modes. $\eta^\pm$ stand for generic rapidity regulators (cut-offs) necessary to separate the soft and collinear modes. The soft function (B1+B2) is split into two pieces along the "separating line" $k^+ = k^-$, allowing for the separation of the whole space into two regions, i.e., two well-defined TMDPDFs: A+B1 (positive rapidities) and C+B2 (negative rapidities). The thick dashed line is the bisectrix of $\tilde{S}(\eta^+,\eta^-)$.*

In Fig. 1 we have denoted by $\eta^\pm$ generic rapidity regulators that separate soft modes from the $n,\bar{n}$-collinear ones. Those regulators should disappear when one combines all the matrix elements within the factorization theorem, since in full QCD there are no rapidity divergencies.

The hadronic tensor for the $q_T$-dependent DY spectrum can be "factorized" [3] in impact parameter space (IPS) as [3]

$$\tilde{M} = H(Q^2)\,\tilde{J}_n^{(0)}(\eta^-)\,\tilde{J}_{\bar{n}}^{(0)}(\eta^+)\,\tilde{S}(\eta^+,\eta^-)\,, \qquad (1)$$

where we show explicitly the rapidity regulator dependence. $\tilde{S}$ is the relevant soft function and $\tilde{J}_{n(\bar{n})}^{(0)}$ stand for *pure* (anti)collinear contributions. Notice that $\tilde{J}_{n(\bar{n})}^{(0)}$ and $\tilde{S}$ will depend also on other parameters that regulate the genuine infra-red (IR) divergencies which exist also in full QCD. Pure collinear matrix elements are calculated by first integrating over all momentum space and then subtracting the "zero-bin" contribution, which is the soft limit of the collinear integrands [4] (see also [15]). Generally speaking, this should be done on a diagram-by-diagram basis. At operator level one can identify those soft contaminations with the soft function itself [16–18], however this equivalence might get spoiled for certain nUV regulators [4].

Thus at the level of the factorization theorem itself, one should refrain from subtracting the soft function (since this would be based on a regulator-dependent arguments) and formulate the relevant theorem in terms of pure collinear matrix elements and soft functions–whenever they are needed–as in Eq. (1).

At the operator level one defines the collinear matrix element as

$$J_n = \frac{1}{2}\sum_{\sigma_1}\langle P\,\sigma_1|\,\overline{\Psi}_n(0^+,y^-,\vec{y}_\perp)\,\frac{\bar{\not{n}}}{2}\,\Psi_n(0)\,|P\,\sigma_1\rangle\ , \qquad (2)$$

where $\Psi_n = W_n^\dagger T_n^\dagger \xi_n$. $W_n$ and $T_n$, collinear and transverse Wilson lines respectively, guarantee the gauge invariance of the matrix element [19, 20]. In the above expression it is implied that the soft contamination has not been subtracted yet. Thus we will refer to it as the "naive" collinear matrix element. The soft function relevant for the $q_T$-dependent DY spectrum is given by

$$S = \langle 0|\,\mathrm{Tr}\,\left[S_n^{T\dagger}S_{\bar{n}}^{T}\right](0^+,0^-,\vec{y}_\perp)\,\left[S_{\bar{n}}^{T\dagger}S_n^{T}\right](0)\,|0\rangle\ , \qquad (3)$$

where $S_n(x) = P\exp\left[ig\int_{-\infty}^{0} ds\, n\cdot A_s(x+sn)\right]$ and the superscript $T$ stand for soft and transverse Wilson lines, respectively. The relevant definitions of the collinear, soft and transverse Wilson lines for DY and DIS kinematics can be found in [3].

---

[3] The quotation marks indicate that Eq. (1) should be considered as an intermediate step towards the final factorization theorem.

[4] Explicit examples can be found in Eq. (64) of [4], and also in [9] where the zero-bin contributions vanish beyond tree-level while the soft function does not.

When the soft function is split symmetrically between $n$ and $\bar{n}$ and combined with the pure collinear matrix elements as in [3], then the TMDPDF defined in this way will depend on the two rapidity regulators $\eta^{\pm}$. The reason is that the soft function is not symmetric with respect to the "separating line" $k^+ = k^-$, which separates negative and positive rapidities in Fig. 1, but it is symmetric with respect to the thick dashed purple line.

To illustrate the above points let us consider the TMDPDF proposed in [3],

$$\tilde{F}_n^{\mathrm{old}} = \tilde{J}_n^{(0)}(\eta^-)\sqrt{\tilde{S}(\eta^+,\eta^-)}\,. \tag{4}$$

The $\Delta$-regulator was used to regularize all nUV divergences, namely the IR and the rapidity divergences. This is why it remains in Eq.(1) when all the matrix elements are combined thus generating the genuine IR of QCD, which in turn is washed out by confinement.

Its $\mathcal{O}(\alpha_s)$ result, before renormalization, is

$$\begin{aligned}\tilde{F}_{n1}^{\mathrm{old}} = \frac{\alpha_s C_F}{2\pi}\Bigg\{&\delta(1-x_n)\left[\frac{1}{\varepsilon_{\mathrm{UV}}^2} - \frac{1}{\varepsilon_{\mathrm{UV}}}\ln\frac{Q^2\Delta^-}{\mu^2\Delta^+} + \frac{3}{2\varepsilon_{\mathrm{UV}}}\right.\\ &\left.-\frac{1}{2}L_T^2 + \frac{3}{2}L_T - L_T\ln\frac{Q^2\Delta^-}{\mu^2\Delta^+} - \frac{\pi^2}{12}\right] + (1-x_n) - L_T\mathcal{P}_{q/q}\\ &-\mathcal{P}_{q/q}\ln\frac{\Delta^-}{\mu^2} - \frac{1}{4}\delta(1-x_n) - (1-x_n)[1+\ln(1-x_n)]\Bigg\}\,,\end{aligned} \tag{5}$$

where $L_T = \ln(\mu^2 b^2 e^{2\gamma_E}/4)$ and $\mathcal{P}_{q/q}$ is the Altarelli-Parisi splitting kernel of a quark in a quark. The existence of two different values for the $\Delta$-regulator is meant to treat in the most general way the supposedly decoupled two collinear sectors. The first line in Eq.(5) contains mixed UV/nUV term. The second line is what would be the matching (or Wilson) coefficient of the TMDPDF onto the integrated PDF after an OPE is carried out, but it also contains unacceptable dependence on the nUV regulator. As it is well-known, Wilson coefficients should be free from any nUV regulators. The last line is simply the integrated PDF. It is clear that only with the choice $\Delta^+ = \Delta^- = \Delta$ the mixed UV/nUV divergence cancels, also the $\Delta$-dependence from the Wilson coefficient. It could be straightforwardly checked that if the latter cancelation does not occur, then one cannot obtain the integrated PDF from the TMDPDF. Thus we could consider the TMDPDF given in Eq. (4) as an acceptable candidate for a TMDPDF *with the prescription of taking the* $\Delta$*'s equal.* This observation was mentioned explicitly in [10]. However in order to avoid such prescription for the regulators a new definition of the TMDPDF should be adopted. We will show below that the proposed new definition will reduce to the one given above when the special case: $\Delta^+ = \Delta^-$ is considered.

It should be immediately mentioned that if one adopts the definition of the TMDPDF given in [3] then the above observation about the (non-)cancellation of UV/nUV divergencies or the $\Delta$-dependence in the Wilson coefficient does *not* rely on the use of the $\Delta$-regulator. In other words it is a feature of the definition rather than the regulators. If one had used off-shellnesses [14], or quark (and gluon) masses, or the $\nu$-regulator as in [9], then the mixed UV/nUV and the nUV regulator dependence could only be canceled when taking the *same* values of the regularizing parameters for both collinear sectors.

However for the general case where $\eta^+ \neq \eta^-$ this is not the case, which sheds doubt on the validity of the definition given in Eq. (4). Thus one should strive to identify the positive rapidity quanta with one collinear sector and the negative rapidity quanta with the other one. This will be achieved by splitting the soft function into two regions, with no overlap between negative and positive rapidity regions (unlike the case of just taking the square root of the soft function and combine it with each of the two pure collinear sectors). We will show next that this splitting is a fundamental property of the soft function, Eq. (3), which holds to all orders in perturbation theory. In our arguments we will refer to the $\Delta$-regulator, however those arguments are independent of the use of any specific regulator.

**Splitting the Soft Function and Definition of TMDPDF:** In the kinematical region where $Q \gg q_T \gg \Lambda_{\mathrm{QCD}}$ one can perform an OPE of the result in Eq. (1) onto the integrated PDFs as [5]

$$\begin{aligned}\tilde{M} = H(Q^2/\mu^2)\,&\tilde{C}(x_n, x_{\bar{n}}; L_T, Q^2/\mu^2)\\ &\times \phi_n(x_n; \Delta^-/\mu^2)\,\phi_{\bar{n}}(x_{\bar{n}}; \Delta^+/\mu^2)\,.\end{aligned} \tag{6}$$

Since the integrated PDFs, $\phi_{n(\bar{n})}$, contain just the $n, \bar{n}$ IR collinear divergences, then with the use of the $\Delta$-regulator each one of the PDFs will depend on a single $\ln\frac{\Delta^-}{\mu^2}$ ($\ln\frac{\Delta^+}{\mu^2}$) to all orders in perturbation theory. Notice also that the matching coefficients $H$ and $\tilde{C}$ cannot depend on any nUV regulator, i.e., they are $\Delta$-free which is also true to all orders in perturbation theory.

On the other hand we can express the hadronic tensor in terms of matrix elements, as in Eq.(1). To separate the modes in rapidity we notice that each one of the pure collinear matrix elements depend just on one regulator, the one which is relevant for their own sector, while the soft function depends on both regulators.

---

[5] $\tilde{C}(x_n, x_{\bar{n}}; L_T, Q^2/\mu^2) = \tilde{C}_n^{\not{Q}}(x_n; L_T)\,\tilde{C}_{\bar{n}}^{\not{Q}}(x_{\bar{n}}; L_T)\,\left(\frac{Q^2}{\mu^2}\right)^{-2D(L_T)}$ in the notation of [3].

Now, given that all the IR of QCD is generated by the two PDFs, as shown in Eq. (6), and that the pure collinear matrix elements depend just on their own regulator, as shown in Eq. (1) combined with the symmetry between $n$ and $\bar{n}$ in the soft function, we immediately derive to all orders in perturbation theory the splitting of the soft function, which in terms of a generic rapidity regulator is

$$\tilde{S}(\eta^+,\eta^-) = \sqrt{\tilde{S}(\eta^+,\eta^+)\,\tilde{S}(\eta^-,\eta^-)}\,. \tag{7}$$

In other words, the soft function $\tilde{S}(\eta^+,\eta^-)$ can be written as the geometric mean of $\tilde{S}(\eta^+,\eta^+)$ and $\tilde{S}(\eta^-,\eta^-)$ [6]. In general, $\tilde{S}(\eta,\eta)$ would be the soft function where we implement the same rapidity regulator for both $n$ and $\bar{n}$ soft Wilson lines. In the case of the $\Delta$-regulator this translates to using the same value of $\Delta^\pm$ for both Wilson lines. For the $\nu$-regulator introduced in [9] it would be to use the same $\nu$. For any other regulator the implementation of Eq. (7) should be straightforward.

From a pictorial point of view, $\tilde{S}(\eta^+,\eta^-)$ would be the area between the lines $\eta^+$ and $\eta^-$ in Fig. 1. Notice that for $\eta^+ \neq \eta^-$ the soft function is not symmetric with respect to the separating line, thus its square root contains both positive and negative rapidity soft quanta. On the other hand, $\tilde{S}(\eta^-,\eta^-)$ would be the area between the two $\eta^-$ lines, symmetric with respect to the separating line. Then, it is clear that its square root is “half” of the region and contains only positive rapidity soft quanta. Thus when we combine $\tilde{J}_n(\eta^-)$ with $\sqrt{\tilde{S}(\eta^-,\eta^-)}$ we collect all the positive rapidity quanta, then defining properly the TMDPDF in the $n$-direction. Same arguments apply for the $\bar{n}$-direction.

Since the soft function can indeed be separated into positive and negative rapidities along the separating line in Fig. 1, we then define the $n$-TMDPDF as

$$\tilde{F}_n = \tilde{J}^{(0)}_n(\eta^-)\sqrt{\tilde{S}(\eta^-,\eta^-)}\,, \tag{8}$$

and for $\tilde{F}_{\bar{n}}$ we interchange $\eta^-$ with $\eta^+$.

The pure collinear matrix element was calculated in [3] with the $\Delta$-regulator,

$$\begin{aligned}\tilde{J}^{(0)}_{n1} = \frac{\alpha_s C_F}{2\pi}\Bigg\{&\delta(1-x_n)\left[\frac{2}{\varepsilon^2_{\rm UV}} - \frac{2}{\varepsilon_{\rm UV}}\ln\frac{\Delta^-}{\mu^2} + \frac{3}{2\varepsilon_{\rm UV}}\right.\\ &\left.-\frac{1}{4} - \frac{2\pi^2}{12} - L_T^2 + \frac{3}{2}L_T - 2L_T\ln\frac{\Delta^-}{\mu^2}\right]\\ &-(1-x_n)\ln(1-x_n) - \mathcal{P}_{q/q}\ln\frac{\Delta^-}{\mu^2} - L_T\mathcal{P}_{q/q}\Bigg\}\end{aligned} \tag{9}$$

which, as explained before, depends just on $\Delta^-$, and the soft function is

$$\begin{aligned}\tilde{S}_1(\Delta^+,\Delta^-) &= \frac{\alpha_s C_F}{2\pi}\left[-\frac{2}{\varepsilon^2_{\rm UV}} + \frac{2}{\varepsilon_{\rm UV}}\ln\frac{\Delta^+\Delta^-}{\mu^2 Q^2}\right.\\ &\qquad\left.+L_T^2 + 2L_T\ln\frac{\Delta^+\Delta^-}{\mu^2 Q^2} + \frac{\pi^2}{6}\right]\\ &= \frac{1}{2}\left[\tilde{S}_1(\Delta^+,\Delta^+) + \tilde{S}_1(\Delta^-,\Delta^-)\right]\,,\end{aligned} \tag{10}$$

thus establishing Eq.(7) at $\mathcal{O}(\alpha_s)$. Combining the pure collinear with $\tilde{S}_1(\Delta^-,\Delta^-)$ that can be extracted from Eq.(10), then the newly defined TMDPDF given in Eq. (8) is

$$\begin{aligned}\tilde{F}_{n1} = \frac{\alpha_s C_F}{2\pi}\Bigg\{&\delta(1-x_n)\left[\frac{1}{\varepsilon^2_{\rm UV}} - \frac{1}{\varepsilon_{\rm UV}}\ln\frac{Q^2}{\mu^2} + \frac{3}{2\varepsilon_{\rm UV}}\right.\\ &\left.-\frac{1}{2}L_T^2 + \frac{3}{2}L_T - L_T\ln\frac{Q^2}{\mu^2} - \frac{\pi^2}{12}\right] + (1-x_n) - L_T\mathcal{P}_{q/q}\\ &-\mathcal{P}_{q/q}\ln\frac{\Delta^-}{\mu^2} - \frac{1}{4}\delta(1-x_n) - (1-x_n)[1+\ln(1-x_n)]\Bigg\}\,.\end{aligned} \tag{11}$$

From the above result we notice that there are no more mixed UV/nUV divergencies as promised, thus it is straightforward to renormalize the TMDPDF. Also the disappearance of the $\Delta$-dependence from the matching coefficient of the TMDPDF onto the integrated PDF (the second line in the previous equation) which allows us to obtain the latter from the former by simple integration in momentum space.

Finally, the fact the TMDPDF is free from the rapidity regulator gives us an evolution equation,

$$\frac{d}{d\eta^-}\tilde{F}_n = 0\,, \tag{12}$$

allowing us to resum large logarithms as it was done in [3]. One more remark is in order. The soft function in Eq.(3) does not have any reference to the high scale $Q^2$, however the $Q^2$-dependence in Eq.(10) is generated by the $\Delta$-regulator in subtle way, because the parameters that regulate the soft Wilson lines, i.e., $\delta^\pm$ (see [3]), are related to $\Delta^\pm$ through the large components of the parton momenta. Although this observation depends on the particular choice of regulators, the fact is that the $Q^2$ in the

---

[6] Eq.(7) can be also derived using equation (10.113) in [8].

TMDPDF is generated by a product of the large components $p^+$ and $\bar{p}^-$, and not $(p^+)^2$ or $(\bar{p}^-)^2$, thus one concludes that the TMDPDF is boost invariant, as it should be. In other words, the TMDPDF can be extracted from a fixed target experiment and used in a collider one, or viceversa.

**Factorization Theorem:** Given the definition of TMDPDF in Eq.(8) we are able to separate positive and negative rapidity modes unambiguously, thus the factorization theorem in Eq.(1), built from ill-defined matrix elements, should be re-written as

$$\tilde{M} = H(Q^2/\mu^2)\, \tilde{F}_n(x_n, b; Q^2, \mu^2)\, \tilde{F}_{\bar{n}}(x_{\bar{n}}, b; Q^2, \mu^2)\,, \quad (13)$$

where the two TMDPDFs are the main hadronic matrix elements. This factorization theorem is valid for $Q \gg q_T \gtrsim \Lambda_{\rm QCD}$, and it can be refactorized in terms of integrated PDFs when $q_T \gg \Lambda_{\rm QCD}$. Notice that this step needs each TMDPDF to be matched independently onto the corresponding PDF, and this is accomplished only if one separates positive and negative rapidity quanta as proposed above. This one-to-one correspondence allows eventually to recover the PDF from the TMDPDF. This passage could not be carried out using Eq. (1) unless the $\Delta$'s are set equal since there is no clean separation of negative and positive rapidities. This requirement can be relaxed with our new definition of the TMDPDF.

**Conclusions and Outlook:** We have provided a definition of a TMDPDF which is free from all rapidity divergencies including the mixed terms that spoil the renormalization of such quantities. This is extremely important for phenomenological applications of TMDs (quarks in the case of DY or SIDIS or gluon TMDPDFs for LHC physics) since the anomalous dimensions and the $Q^2$-resummation kernel $D$ [3], and thus the evolution of individual TMDs can be unambiguously determined in the sense that no un-physical extra parameters are needed. Moreover, the resummation of all large logarithms can be obtained by “running” between the different relevant scales and by solving a homogeneous evolution equation as in Eq. (12). Our results can be readily extended to define polarized and unpolarized quark and gluon TMDPDFs which could be considered as a generalization of the one introduced in [21], as well as TMD fragmentation functions. The definition of TMDPDF given in this letter reduces to the one proposed in [3] when the two sectors are treated identically. We finally remark that our treatment of the TMDPDF applies for massless and massive partons as well, where the quark masses combined with an artificial gluon mass serve as IR regulators, but not rapidity regulators; for the latter one needs an additional set of regulators.

*Acknowledgements:* This work is supported by the Spanish MEC, FPA2011-27853-CO2-02. M.G.E. is supported by the PhD funding program of the Basque Country Government. M.G.E. thanks LBNL for its hospitality while part of this work was completed. A.I. would like to thank Andreas Schäfer for useful discussions. I.S. is supported by the Ramón y Cajal Program.

---

[*] Electronic address: miguel.gechevarria@fis.ucm.es
[†] Electronic address: idilbi@ectstar.eu
[‡] Electronic address: ignazios@fis.ucm.es

[1] C. W. Bauer, D. Pirjol, and I. W. Stewart, Phys.Rev. **D67**, 071502 (2003), hep-ph/0211069.

[2] S. Mantry and F. Petriello, Phys.Rev. **D81**, 093007 (2010), 0911.4135.

[3] M. G. Echevarria, A. Idilbi, and I. Scimemi, JHEP **1207**, 002 (2012), 1111.4996.

[4] A. V. Manohar and I. W. Stewart, Phys.Rev. **D76**, 074002 (2007), hep-ph/0605001.

[5] J. C. Collins and D. E. Soper, Nucl.Phys. **B194**, 445 (1982).

[6] J. C. Collins and D. E. Soper, Nucl.Phys. **B193**, 381 (1981).

[7] X.-d. Ji, J.-p. Ma, and F. Yuan, Phys.Rev. **D71**, 034005 (2005), hep-ph/0404183.

[8] J. Collins, *Foundations of perturbative QCD* (Cambridge monographs on particle physics, nuclear physics and cosmology. 32, 2011).

[9] J.-Y. Chiu, A. Jain, D. Neill, and I. Z. Rothstein, JHEP **1205**, 084 (2012), 1202.0814.

[10] M. G. Echevarria, A. Idilbi, and I. Scimemi (2012), 1209.3892.

[11] I. Cherednikov and N. Stefanis, Nucl.Phys. **B802**, 146 (2008), 0802.2821.

[12] J. C. Collins and T. C. Rogers (2012), 1210.2100.

[13] T. Becher and M. Neubert, Eur.Phys.J. **C71**, 1665 (2011), 1007.4005.

[14] J. Chay and C. Kim (2012), 1208.0662.

[15] J.-y. Chiu, A. Fuhrer, A. H. Hoang, R. Kelley, and A. V. Manohar, Phys.Rev. **D79**, 053007 (2009), 0901.1332.

[16] C. Lee and G. F. Sterman, Phys.Rev. **D75**, 014022 (2007), hep-ph/0611061.

[17] A. Idilbi and T. Mehen, Phys.Rev. **D75**, 114017 (2007), hep-ph/0702022.

[18] A. Idilbi and T. Mehen, Phys.Rev. **D76**, 094015 (2007), 0707.1101.

[19] M. Garcia-Echevarria, A. Idilbi, and I. Scimemi, Phys.Rev. **D84**, 011502 (2011), 1104.0686.

[20] A. Idilbi and I. Scimemi, Phys.Lett. **B695**, 463 (2011), 1009.2776.

[21] M. G. Echevarria, A. Idilbi, A. Schafer, and I. Scimemi (2012), 1208.1281.